



Published in final edited form as:

Tetrahedron Lett. 2015 August 5; 56(32): 4670–4673. doi:10.1016/j.tetlet.2015.06.039.

Synthesis and Exploration of Electronically Modified (*R*)-5,5-Dimethyl-(*p*-CF₃)₃-*i*-PrPHOX in Palladium-Catalyzed Enantio- and Diastereoselective Allylic Alkylation: A Practical Alternative to (*R*)-(*p*-CF₃)₃-*t*-BuPHOX


Robert A. Craig II and Brian M. Stoltz*

Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA 91125, United States

Abstract

The synthesis of the novel electronically modified phosphinooxazoline (PHOX) ligand, (*R*)-5,5-dimethyl-(*p*-CF₃)₃-*i*-PrPHOX, is described. The utility of this PHOX ligand is explored in both enantio- and diastereoselective palladium-catalyzed allylic alkylations. These investigations prove (*R*)-5,5-dimethyl-(*p*-CF₃)₃-*i*-PrPHOX to be an effective and cost-efficient alternative to electronically modified PHOX ligands derived from the prohibitively expensive (*R*)-*t*-leucine.

Graphical abstract



- **Cost-Effective** Alternative to *t*-BuPHOX
- **Electronically Modified** Architecture
- **Highly Effective** in Enantio- and Diastereoselective Palladium-Catalyzed Allylic Alkylation

Keywords

Allylic Alkylation; Diastereoselective; Enantioselective; Palladium-catalyzed; Phosphinooxazoline

*Corresponding author. Tel.: +1-626-395-6064; fax: +1-626-395-8436; stoltz@caltech.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary data

NMR spectra for new compounds (i.e., **13**, **14**, and (*R*)-**L5**) can be found in the supporting information, which is available online at: <http://>

1. Introduction

Phosphinooxazoline (PHOX) ligands, developed by Helmchen,¹ Williams,² and Pfaltz,³ have proven to be a privileged ligand scaffold in transition metal catalysis.⁴ PHOX ligands have found application in a variety of asymmetric transition metal-catalyzed transformations including asymmetric hydrogenation,⁵ azomethine ylide cycloadditions,⁶ intermolecular Heck couplings,⁷ and hydrosilylation⁸ as well as transition metal-catalyzed allylic substitution^{4,9} and protonation¹⁰ reactions. Our lab has extensively explored the utility of the PHOX ligand scaffold in the palladium-catalyzed enantioselective allylic alkylation of carbocyclic¹¹ and heterocyclic¹² substrates. These investigations have revealed electronically modified PHOX ligands (i.e. (*S*)-(p-CF₃)₃-*t*-BuPHOX ((*S*)-**L1**), Figure 1)¹³ can profoundly enhance the rate of reaction as well as yield, enantiomeric excess (*ee*) and/or diastereomeric ratio of a product containing an all-carbon quaternary center (e.g. use of (*S*)-**L1** vs. (*S*)-**L2** to construct lactam **2**,^{12e} cyclohexanone **4**,^{13c} cyclohexenone **6**,^{13b} and cyclohexanone diastereomers **9** and **10**,¹⁴ Schemes 1A–1C and Scheme 2, respectively).

Most commonly, transition metal complexes employing *tert*-leucinol-derived PHOX ligands (e.g. (*S*)-**L1** and (*S*)-**L2**, Figure 1) enable the formation of the corresponding products with the best enantiomeric and diastereomeric ratios. Although (*R*)-*t*-BuPHOX has been employed in natural product synthesis¹⁵ and explored in transition-metal catalyzed allylic alkylations,^{10a,16} these examples are quite rare considering the nearly prohibitive cost of the requisite starting material for ligand synthesis, (*R*)-*t*-leucine.¹⁷ Previously, 5,5-geminally disubstituted (*R*)-valine-derived PHOX ligands (e.g. (*R*)-**L3** and (*R*)-**L4**, Figure 2) have been constructed as cost-effective alternatives to (*R*)-*t*-BuPHOX ((*R*)-**L2**).¹⁸ We sought to extend this precedent to the synthesis of electronically modified congener (*R*)-5,5-dimethyl-(p-CF₃)₃-*i*-PrPHOX ((*R*)-(p-CF₃)₃-*i*-PrPHOX^{Me2}, (*R*)-**L5**, Figure 2) and explore its efficacy as a ligand in palladium-catalyzed enantio- and diastereoselective allylic alkylation reactions.

2. Results and discussion

2.1 Synthesis of (*R*)-(p-CF₃)₃-*i*-PrPHOX^{Me2} ((*R*)-**L5**)

Synthesis of (*R*)-(p-CF₃)₃-*i*-PrPHOX^{Me2} ((*R*)-**L5**) was initiated with acid chloride **11**¹⁹ and the hydrogen chloride salt of (*R*)-valine derivative **12**¹⁸ (Scheme 3). Intermolecular coupling of acid chloride **11** and amino alcohol **12** in the presence of excess Et₃N provides amide **13** in 79% yield. Intramolecular cyclization of amide **13** under acidic conditions furnishes oxazoline **14** in 87% yield. Completion of desired ligand (*R*)-**L5** was accomplished over two steps, beginning with the copper-mediated coupling of phosphine oxide **15** with bromide **14** at elevated temperature.²⁰ This procedure produces phosphine oxide **16** in 63% yield. Reduction of phosphine oxide **16** was subsequently accomplished in neat Ph₂SiH₂ at 140 °C over 48 hours, providing the desired ligand (*R*)-(p-CF₃)₃-*i*-PrPHOX^{Me2} ((*R*)-**L5**) in 81% yield in the final step of the synthetic sequence.

2.2 Use of (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} in Palladium-Catalyzed Asymmetric Transformations

Application of (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) was initially explored in the intermolecular palladium-catalyzed enantioselective allylic alkylation of silyl enol ether **17** with mesylate **18** (Scheme 4). Previously we disclosed the initial development and optimization of this transformation using (*S*)-*t*-BuPHOX (**(S)-L2**), which afforded chloroallylketone (**(S)-19**) in 82% yield and 92% *ee* (entry 1).^{12d} Substitution of (**(S)-L2**) with the electronically modified (*S*)-(p-CF₃)₃-*t*-BuPHOX (**(S)-L1**) provided the product (**(S)-19**) in a slightly diminished 91% *ee* (entry 2). Switching the ligand to (*S*)-5,5-diphenyl-*i*-PrPHOX (**(S)-L3**) furnished chloroallylketone (**(S)-19**) in 90% *ee* (entry 3). Moving into the opposite enantiomeric series, the use of (*R*)-5,5-dimethyl-*i*-PrPHOX (**(R)-L4**) provided chloroallylketone (**(R)-19**) in a somewhat diminished 89% *ee* (entry 4) compared to the originally optimized reaction conditions (entry 1). Alternatively, we were pleased to find that (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) afforded chloroallylketone (**(R)-19**) in the same 91% *ee* (entry 5) in the opposite enantiomeric series compared to the use of (*S*)-(p-CF₃)₃-*t*-BuPHOX (entry 2). It is noteworthy that the required reaction time and isolated yield of chloroallylketone **19** were independent of the ligand employed. Thus, (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) can allow access to the enantiomeric series of products to those afforded in reactions employing (**(S)-L1**) without any loss in product *ee* in a cost-effective manner, being derived from (*R*)-valine, which is less than 2% of the cost of (*R*)-*t*-leucine.

The utility of (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) was further demonstrated in the intermolecular palladium-catalyzed diastereoselective decarboxylative allylic alkylation of β-ketoester **20** with allyl electrophile **21** (Scheme 5).^{16a} While the system displays an inherent selectivity for the formation of diastereomer **22** in a 2:1 ratio with diastereomer **23** when achiral PHOX ligand **L6** was employed (entry 1),²¹ the use of (*S*)-*t*-BuPHOX (**(S)-L2**) can override this substrate bias, providing diastereomer **23** as the major product (entry 2). Comparatively, the use of (*R*)-*t*-BuPHOX (**(R)-L2**) reinforces the inherent selectivity, providing diastereomer **22** in a 12:1 ratio with minor diastereomer **23** in a combined 73% yield (entry 3). Pleasingly, the employment of (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) further improved this transformation, furnishing an 18:1 mixture of products in favor of diastereomer **22** in an improved 85% combined yield (entry 4). These studies revealed that (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) was the optimal ligand for the highly diastereoselective formation of allylic alkylation product **22**. Additionally, other research groups have found (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} (**(R)-L5**) to be a uniquely effective ligand for the palladium-catalyzed diastereoselective allylic alkylation of other carbocyclic substrates.²²

3. Conclusion

Herein, we have disclosed the synthesis of a new, electronically modified phosphinooxazoline (PHOX) ligand, (*R*)-5,5-dimethyl-(p-CF₃)₃-i-PrPHOX (**(R)-(p-CF₃)₃-i-PrPHOX^{Me2}**, (**(R)-L5**). Derived from (*R*)-valine, this cost-effective alternative to (*R*)-(p-CF₃)₃-*t*-BuPHOX (**(R)-L1**) has proved effective in both palladium-catalyzed enantio- and diastereoselective allylic alkylations, furnishing the alkylation products in comparable *ee* and improved diastereomeric ratio. Efforts to further explore the utility of the readily

available (*R*)-(p-CF₃)₃-i-PrPHOX^{Me2} ligand in palladium-catalyzed stereoselective transformations are currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors wish to thank the NIH-NIGMS (R01GM080269), Amgen, the Gordon and Betty Moore Foundation, and Caltech for financial support. R.A.C. gratefully acknowledges the support of this work provided by a fellowship from the National Cancer Institute of the National Institutes of Health under Award Number F31A17435.

References

1. Sprinz J, Helmchen G. *Tetrahedron Lett.* 1993; 34:1769–1772.
2. Dawson GJ, Frost CG, Williams MJ, Coote SJ. *Tetrahedron Lett.* 1993; 34:3149–3150.
3. Von Matt P, Pfaltz A. *Angew. Chem. Int. Ed. Engl.* 1993; 32:566–568.
4. a Carroll MP, Guiry PJ. *Chem. Soc. Rev.* 2014; 43:819–833. [PubMed: 24257099] b Hargaden GC, Guiry PJ. *Chem. Rev.* 2009; 109:2505–2550. [PubMed: 19378971] c McManus HA, Guiry PJ. *Chem. Rev.* 2004; 104:4151–4202. [PubMed: 15352789] d Helmchen G, Pfaltz A. *Acc. Chem. Res.* 2000; 33:336–345. [PubMed: 10891051] e Williams JM. *Synlett.* 1996:705–710.
5. a Verendel JJ, Pàmies O, Diéguez M, Andersson PG. *Chem. Rev.* 2014; 114:2130–2169. [PubMed: 24568181] b Braunstein P, Graiff C, Naud F, Pfaltz A, Tiripicchio A. *Inorg. Chem.* 2000; 39:4468–4475.
6. Stohler R, Wahl F, Pfaltz A. *Synthesis.* 2005:1431–1436.
7. a Gilbertson SR, Fu Z. *Org. Lett.* 2001; 3:161–164. [PubMed: 11430024] b Hashimoto Y, Horie Y, Hayashi M, Saigo K. *Tetrahedron: Asymmetry.* 2000; 11:2205–2210.
8. a Frölander A, Moberg C. *Org. Lett.* 2007; 9:1371–1374. [PubMed: 17323963] b Sudo A, Yoshida H, Saigo K. *Tetrahedron: Asymmetry.* 1997; 8:3205–3208.
9. a Liu Y, Liniger M, McFadden RM, Roizen JL, Malette J, Reeves CM, Behenna DC, Seto M, Kim J, Mohr JT, Virgil SC, Stoltz BM. *Beilstein J. Org. Chem.* 2014; 10:2501–2512. [PubMed: 25383121] b Behenna DC, Mohr JT, Sherden NH, Marinescu SC, Harned AM, Tani K, Seto M, Ma S, Novák Z, Krout MR, McFadden RM, Roizen JL, Enquist JA Jr, White DE, Levine SR, Petrova KV, Iwashita A, Virgil SC, Stoltz BM. *Chem. Eur. J.* 2011; 17:14199–14223. [PubMed: 22083969] c García-Yebra C, Janssen JP, Rominger F, Helmchen G. *Organometallics.* 2004; 23:5459–5470.
10. a Doran R, Carroll MP, Akula R, Hogan BF, Martins M, Fanning S, Guiry PJ. *Chem. Eur. J.* 2014; 20:15354–15359. [PubMed: 25314579] b Carroll MP, Müller-Bunz H, Guiry PJ. *Chem. Commun.* 2012; 48:11142–11144. c Marinescu SC, Nishimata T, Mohr JT, Stoltz BM. *Org. Lett.* 2008; 10:1039–1042. [PubMed: 18303896] d Mohr JT, Nishimata T, Behenna DC, Stoltz BM. *J. Am. Chem. Soc.* 2006; 128:11348–11349. [PubMed: 16939246]
11. a Numajiri Y, Pritchett BP, Chiyoda K, Stoltz BM. *J. Am. Chem. Soc.* 2015; 137:1040–1043. [PubMed: 25578104] b Reeves CM, Behenna DC, Stoltz BM. *Org. Lett.* 2014; 16:2314–2317. [PubMed: 24724872] c Reeves CM, Eidamshaus C, Kim J, Stoltz BM. *Angew. Chem. Int. Ed.* 2013; 52:6718–6721. d Enquist JA Jr, Stoltz BM. *Nature.* 2008; 453:1228–1231. [PubMed: 18580947] e Mohr JT, Behenna DC, Harned AM, Stoltz BM. *Angew. Chem. Int. Ed.* 2005; 44:6924–6927. f Behenna DC, Stoltz BM. *J. Am. Chem. Soc.* 2004; 126:15044–15045. [PubMed: 15547998]
12. a Numajiri Y, Jiménez-Osés G, Wang B, Houk KN, Stoltz BM. *Org. Lett.* 2015; 17:1082–1085. [PubMed: 25714704] b Korch KM, Eidamshaus C, Behenna DC, Nam S, Horne D, Stoltz BM. *Angew. Chem. Int. Ed.* 2015; 54:179–183. c Bennett NB, Duquette DC, Kim J, Liu W-B, Marziale AN, Behenna DC, Virgil SC, Stoltz BM. *Chem. Eur. J.* 2013; 19:4414–4418. [PubMed: 23447555] d Craig RA II, Roizen JL, Smith RC, Jones AC, Stoltz BM. *Org. Lett.* 2012; 14:5716–

5719. [PubMed: 23101616] e Behenna DC, Liu Y, Yurino T, Kim J, White DE, Virgil SC, Stoltz BM. *Nature Chem.* 2012; 4:130–133. [PubMed: 22270628] f Seto M, Roizen JL, Stoltz BM. *Angew. Chem. Int. Ed.* 2008; 47:6873–6876.
13. a McDougal NT, Streuff J, Mukherjee H, Virgil SC, Stoltz BM. *Tetrahedron Lett.* 2010; 51:5550–5554. [PubMed: 21076623] b White DE, Stewart IC, Grubbs RH, Stoltz BM. *J. Am. Chem. Soc.* 2008; 130:810–811. [PubMed: 18163634] c Tani K, Behenna DC, McFadden RM, Stoltz BM. *Org. Lett.* 2007; 9:2529–2531. [PubMed: 17536810]
14. Streuff J, White DE, Virgil SC, Stoltz BM. *Nature Chem.* 2010; 2:192–196. [PubMed: 20697457]
15. a Day JJ, McFadden RM, Virgil SC, Kolding H, Alleva JL, Stoltz BM. *Angew. Chem. Int. Ed.* 2011; 50:6814–6818. b Levine SR, Krout MR, Stoltz BM. *Org. Lett.* 2009; 11:289–292. [PubMed: 19093809] c Petrova KV, Mohr JT, Stoltz BM. *Org. Lett.* 2009; 11:293–295. [PubMed: 19093836]
16. a Liu W-B, Reeves CM, Virgil SC, Stoltz BM. *J. Am. Chem. Soc.* 2013; 135:10626–10629. [PubMed: 23829704] b Fang X, Johannsen M, Yao S, Gathergood N, Hazell RG, Jørgensen KA. *J. Org. Chem.* 1999; 64:4844–4849. [PubMed: 11674560]
17. The cost of (*R*)-*t*-leucine ranges between \$350 and \$400 per gram, depending on the size of the order from Sigma-Aldrich, as advertised on their sigmaaldrich.com, accessed 30 April, 2015. The synthesis of *t*-BuPHOX ligands, however, can be accomplished with ease on large scale, Mohr JT, Krout MR, Stoltz BM. *Org. Synth.* 2009; 86:194–211. [PubMed: 21197146]
18. a Bélanger é, Pouliot M-F, Courtemanche M-A, Paquin J-F. *J. Org. Chem.* 2012; 77:317–331. [PubMed: 22148548] b Bélanger é, Pouliot M-F, Paquin J-F. *Org. Lett.* 2009; 11:2201–2204. [PubMed: 19388656]
19. Acid chloride **11** was synthesized in two steps from 2-bromo-5-(trifluoromethyl)benzonitrile by a known procedure, see: reference 13b.
20. The procedure for the coupling of phosphine oxide **15** with oxazoline **14** and sequential reduction was adapted from reference 13a.
21. Control experiments were performed using achiral PHOX ligand **L6**, bearing no substituent on the oxazoline ring, see reference 16a for full details.
22. Professor Stephen F. Martin, University of Texas at Austin, personal communication.







Figure 1.
Electronically Modified and Unmodified (S)-*t*-BuPHOX Ligands




KNOWN 5,5-DISUBSTITUTED *i*-PrPHOX LIGANDS



**TARGETED 5,5-DISUBSTITUTED,
ELECTRONICALLY
MODIFIED *i*-PrPHOX LIGAND**


Figure 2.
5,5-Geminally Disubstituted (*R*)-Valine-Derived PHOX ligands

A.




Stoltz, et. al. REFERENCE 12(e)

B.



Stoltz, et. al. REFERENCE 13(c)


C.




Stoltz, et. al. REFERENCE 13(b)

Scheme 1.


Comparison of Electronically Modified (*S*)-(p-CF₃)₃-*t*-BuPHOX (**(S)-L1**) and Unmodified (*S*)-*t*-BuPHOX (**(S)-L2**) in Intramolecular Palladium-Catalyzed Enantioselective Allylic Alkylation

**Scheme 2.**

Comparison of Electronically Modified (*S*)-(p-CF₃)₃-*t*-BuPHOX (**(S)-L1**) and Unmodified (*S*)-*t*-BuPHOX (**(S)-L2**) in Diastereoselective Decarboxylative Alkylation Cascade



Scheme 3.
Synthesis of (R)-(p-CF₃)₃-i-PrPHOX^{Me}₂ ((R)-L5)




Entry	Ligand	Amino Acid Ligand Precursor	Amino Acid Cost per gram (\$) ^a		Product ee (%) ^b
			S - Enantiomer	- R	
1	(S)-L2	<i>tert</i> -Leucine	33	351	−92
2	(S)-L1	<i>tert</i> -Leucine	33	351	−91
3	(S)-L3	Valine	0.60	6	−90
4	(R)-L4	Valine	0.60	6	89
5	(R)-L5	Valine	0.60	6	91

^a Cost per gram of amino acid from Sigma-Aldrich, accessed 4/30/2015.


^b Enantiomeric excess (ee) measured by analytical chiral GC.

Scheme 4.

Ligand Comparison in Enantioselective Palladium-Catalyzed Intermolecular Allylic Alkylation.



Entry	Ligand	% yield ^a	dr (22 : 23) ^b
1	L6	79	2:1
2	(<i>S</i>)- L2	79	1:2
3	(<i>R</i>)- L2	73	12:1
4	(<i>R</i>)- L5	85	18:1



^a Isolated yield of **22** and **23**. ^b Determined by ^1H NMR analysis of the crude reaction mixture and analytical GC analysis

Scheme 5.

Diastereoselective Decarboxylative Allylic Alkylation Employing (*R*)-(*p*- CF_3)₃-*i*-PrPHOX^{Me2} ((*R*)-**L5**).